Minsk 2017

XXIV NPCS MINSK-SOSNY 16-19MAY 2017

TOWARDS THEORY OF THE NEW SPIN STRUCTURES

Sergey Vladimirovich Smurov
Alexandr Albertovich Maslikov
(Engineering Physics Institute, Serpukhov)
Gennady Germanovich Volkov
Peterburg Institute of Nuclear Physics, Gatchina,
Engineering Physics Institute, Serpukhov)

Minsk
2017

Tre-Neutrino Sull~Estensione Multidimensionale Del Continuum Spacio-Temporale Del`Universo Visible

MATTER IN VISIBLE IN UNIVERSE

ФЕРМИОНЫ				БОЗОНЫ	
ДИРАК-ФЕРМИОНЫ	СПИН J=1/2			БОЗОНЫ	СПИН J=1
ЛЕПТОНЫ	e	μ	τ	ФОТОН	A
ДИРАК-ФЕРМИОНЫ	СПИН J=1/2			БОЗОНЫ	СПИН J= 1
КВАРКИ	(u, d)	(c, s)	(t, b)	ГЛЮОНЫ	G=1,..8
	(u, d)	(c, s)	(t, b)		
	(u, d)	(c, s)	(t, b)		
МАЙОРАНО-ФЕРМИОНЫ	СПИН J=1/2			БОЗОНЫ	СПИН J= 1
НЕЙТРИНО	$\mathcal{N}(e)$	$\mathcal{N}(\mu)$	$\mathcal{N}(\tau)$	Weak	W(+),W(-),Z

1. Standard Model and new space-time geometrical structure of the Universe

- The geometrical basis of the modern quantum field theory sucessufully describing the
- $U(1)EM$- electrodynamic processes, the $SU(3c)$-gauge quantum chromodynamics and the
- electroweak interactions based on the $SU(2)WI \times U(1)Y$ - gauge broken symmetry is our
- space-time world what can be represented as a homogeneous and isotropic $D = (3 + 1)$-
- four-dimensional continuum. The symmetry properties of the spatial and temporal continuum
- describe by the Lorentz-Poincar´*e* groups and its representations and some fundamental
- discrete symmetries- P,T,C.

Space-time geometrical structure of the Universe

- This space-time continuum can be immersed into
- much huge comprehensive multidimensional world.
- The modern experimental data derived
- from the elementary particle physics and astrophysics allow us to estimate the sizes
- of the expanding visible part of the continuum
- $\Lambda_{min} \leq \Lambda \leq \Lambda_{max}$.

THE DOWN-UP QUARK MASSES DEPEND ON THE E-M CHARGE AND ON THE NUMBER OF GENERATIONS (NEW CHARGE - ORIGIN FROM D=6?)

$$
\begin{aligned}
m_{i_k} &\approx (q^u)^{2k} m_0, \qquad k = 0,1,2; \qquad i_0 = u,\ i_1 = c,\ i_2 = t, \\
m_{i_k} &\approx (q^d)^{2k} m_0, \qquad k = 0,1,2; \qquad i_0 = d,\ i_1 = s,\ i_2 = b,
\end{aligned}
\tag{1}
$$

where $q^u = (q^d)^2$, $q^d \approx 4 - 5 \approx 1/\lambda$, $\lambda = \sin\theta_C$.

THE FERMIONS MASSES AND W-Z- BOSONS COULD DEFINED BY THE E-W SCALE

M- EW-SCALE THE PHASE TRANSITION BETWEEN TWO VACUUMS???

СВОЙСТВА НЕЙТРИНО О МНОГОМЕРНОМ ОБОБЩЕНИИ ТЕОРИИ ОТНОСИТЕЛЬНОСТИ

ПРЕДСТАВЛЕНИЯ ГРУППЫ ЛОРЕНЦ- ПУАНКАРЕ

- *1. Спин s=½*
- ***2. Майорано-Вейлевская природа** - 2-х компклексные степени свободы*

3. Масса m =O(eV)

ЭЛЕКТРОМАГНИТНАЯ СТЕРИЛЬНОСТЬ

- *4. Заряд Q (EM)= O*
- *5. Магнитный момент Mag=O(0)*

(V – A)- СЛАБАЯ СВЯЗЬ С ЗЛЕКТРОМАГНИТНЫМ МИРОМ

- ***6. Взаимодействие слабое***

1967 ---------- 2017(STO-M(1,3))

- WS-SU(2)XU(1)-Model
- SU(5) and SO(10)—GUT
- Strings + Superstrings
- M11- Superrgravity+Kaluza-Klein Compactifications
- Heterotic SuperstringsE(8)XE(8) Models and K6=CY_3- compactifications
- 4-dim SS with WS Fermions
- D-Membranes
- M11, M12 – and String Duality

ПУТИ РАСШИРЕНИЯ МЕТРИКИ

- **A) стандартный**
- Ds^2=dx_0^2-dx_1^2-dx_2-…-dx_n^2-.....
- Lie algebras and groups SO(p,q)n=p+q – space-time groups and double covered Spin(p,q),…
- **B)Non-standard ways…..T_mnk…**
- New symmetries ----->new groups and algebras, theory of new numbers…
- New geometry- BCY_n, Group algebra Spaces,…

1 Theories of Numbers in Geometry and in Physics.I

The further progress in modern models of elementary particles and cosmology is related to the searching for new Riemann and tensor structures in multidimensional spaces $D \geq 4$ based on the theories of new hyper-numbers, new algebras and new symmetries. The traditional geometry of Riemann and pseudo-Riemann symmetric homogeneous compact and non-compact spaces was associated to the classification of the Killing-Cartan-Lie algebras, according to the theories of binary hyper-numbers,- the well-known real-$\mathbb{R}$, complex- $\mathbb{C}$, quaternions- $\mathbb{H}$, octonions-$\mathbb{O}$- normed division algebras.

TOWARDS A N-ary MATHEMATICS+PHYSICS

- THE WAYS TO EXTRA WORLD
- 1)BCY- SU(n), G2 - Holonomy Geometry
- 2)Theories of the Cyclic C_n- Complex Numbers
- 3)Finite Group Algebras
- **MASS CHARGE SPIN …???**

9 The n-ary Algebra of Reflexive Projective Numbers and CY-classification

A CY_n space can be realized as an algebraic variety $\mathcal{M}$ in a weighted projective space $\mathrm{CP}^{n-1}(\overrightarrow{k})$ where the weight vector reads $\overrightarrow{k} = (k_1, \ldots, k_n)$. This variety is defined by

$$\mathcal{M} \equiv (\{x_1, \ldots, x_n\} \in \mathrm{CP}^{n-1}(\overrightarrow{k}) : \mathcal{P}(x_1, \ldots, x_n) \equiv \sum_{\vec{m}} c_{\vec{m}} x^{\vec{m}} = 0), \tag{6}$$

i.e., as the zero locus of a quasi–homogeneous polynomial of degree

$$d_k = \sum_{i=1}^{n} k_i, \tag{7}$$

with the monomials being

$$x^{\vec{m}} \equiv x_1^{m_1} \cdots x_n^{m_n}. \tag{8}$$

6 Berger classification of non-symmetric Riemann spaces

Firstly, in 1955, Berger presented the classification of irreducibly acting matrix Lie groups occur as the holonomy of a torsion free affine connection. The Berger list of non-symmetric irreducible Riemann manifolds with the list of holonomy groups H of M one can see.

Table 2: *The list of Berger classification for non-symmetric Riemannian spaces.*

M	G_{ISOM}	H_{Hol}	Dim_R	*metrics*
General	–	$SO(n)$	n	
Kahler	–	$U(n) \subset O(2n)$	n	
Calabi – Yau	–	$SU(n) \subset SO(2n)$	$2n$	
Hyper – Kahler	–	$Sp(n) \subset SO(4n)$	$4n$	
quaternion – –Kahler	–	$Sp(n) \times Sp(1) \subset SO(4n)$	$4n$	
exceptional	–	$G(2) \subset SO(7)$	7	
exceptional	–	$spin(7) \subset SO(8)$	8	

BERGER-CALABI-YAU SPACES

HODGE DIAMOND

with

HOLONOMY (g) = SU(n)



$$g = (|dz_1|^2 + \dots\dots + |dz_n|^2)$$

Kahler form

$$\omega = \frac{i}{2}(dz_1 \wedge d\bar{z}_1 + \dots + dz_n \wedge d\bar{z}_n)$$

(n,0) parallel form

$$\Omega = dz_1 \wedge \dots\dots \wedge dz_n$$

$$b_{n0} = b_{0n} = 1$$

$$b_{p0} = b_{0p} = 0$$

$$n > p > 0$$

$$\frac{\omega^n}{n!} = (-1)^{\frac{n(n-1)}{2}} \left(\frac{i}{2}\right)^n \Omega \wedge \bar{\Omega}$$

K3-Manifolds (BCY_2)



G	k_3^{ext}	L ■	J_n ⬢	R ●	k_3^{ext}	G
cyclic	(0,0,0,1)	A_r^1		A_r^1	(0,0,0,1)	cyclic
dihedral	(0,0,1,1)	D_r^1		D_r^1	(0,0,1,1)	dihedral
tetrahedral	(0,1,1,1)	E_6^1		E_6^1	(0,1,1,1)	tetrahedral
octahedral	(0,1,1,2)	E_7^1		E_7^1	(0,1,1,2)	octahedral
icosahedral	(0,1,2,3)	E_8^1		E_8^1	(0,1,2,3)	icosahedral

CY3-Newton polyhedron k=(11248)



K3 polyhedron

(1,0,2,3) + (0,1,2,3) = (1,1,4,6)[12]



Weierstrass triangle

$E(8)^{(1)}$ diagram

$E(8)^{(1)}$ diagram

Cartan–Lie Dynkin diagram



Affine Kac–Moody diagram





Affine binary–ternary Berger diagram

МНОГОМЕРНОЕ РАСШИРЕНИЕ СПЕЦИАЛЬНОЙ ТЕОРИИ ОТНОСИТЕЛЬНОСТИ

- 1Принцип максимальности скорости света будет справедлив только для заряженного вакуума то есть для частиц обладающих электромагнитным зарядом Темная материя и стерильное нейтрино Могут распространяться с гораздо большими скоростями
- 2 Многомерное обобщение группы Лоренца предполагает существование другого буста и возможного раширения понятия времени даже за счет структуры
- 3 принцип относительности также может потребовать расширения

За счет появления новых некомпактифицированых размерностей

стрелки времени или стрелки пространства. Поэтому появляются несколько возможностей поиска параметра энергии “ветровой”или “температурной”, от которой может зависеть скорость нейтрино и мы привели две схемы экспериментов- это должны решить будущие эксперименты

Cartan–Lie Dynkin diagram



Affine Kac–Moody Dynkin diagram





Affine binary–ternary Berger diagram

Cartan–Lie Dynkin diagram



Affine Kac–Moody Dynkin diagram





Affine binary–ternary Berger diagram

COMPLEXIFICATION OF R^n

We will discuss the following themes:

- C^n- hyper-plural division numbers
- C^n- complexification of R^n- spaces, $n = 3, 4, 5, 6, ...,$
- C^n- structure and the invariant surfaces, $n = 3, 4, 5, 6, ...$
- C^n- hyper-holomorphic and hyper-harmonic functions
- The link between C^n-holomorphism and the origin of n-spinors

GEOMETRY OF BINARY HYPER NUMBERS

$$x^2 = 1, x \in R, \tag{44}$$

$$|\hat{Z}| = x_0^2 + x_1^2 = 1 \in \mathbb{C}, \tag{45}$$

$$|\hat{Q}| = x_0^2 + x_1^2 + x_2^2 + x_3^2 = |Z_1|^2 + |Z_2|^2 = 1 \in \mathbb{H}, \tag{46}$$

$$|\hat{O}| = x_0^2 + x_1^2 + \ldots + x_7^2 = |Q_1|^2 + |Q|^2 = 1 \in \mathbb{O}, \tag{47}$$

$$\tag{48}$$

N-ARY HYPER NUMBERS

If the binary alternative division algebras (real numbers, complex numbers, quaternions, octonions) over the real numbers have the dimensions 2^p, $p = 0, 1, 2, 3, 4, ...$, the n=3-ary and n=4-ary norm division algebras have the following dimensions n^p, $p = 0, 1, 2, 3$, respectively:

$$
\begin{array}{ll|ll}
\mathbb{R}: & 2^0 = 1 & \mathbb{R}: & 3^0 = 1 \\
\mathbb{C}: & 2^1 = 1+1 & \mathbb{TC}: & 3^1 = 1+1+1 \\
\mathbb{Q}: & 2^2 = 1+2+1 & \mathbb{TQ}: & 3^2 = 1+2+3+2+1 \\
\mathbb{O}: & 2^3 = 1+3+3+1 & \mathbb{TO}: & 3^3 = 1+3+6+7+6+3+1 \\
\mathbb{S}: & 2^4 = 1+4+6+4+1 & \mathbb{TS}: & 3^4 = 1+4+10+16+19+16+10+4+1
\end{array}
\tag{49}
$$

$$
\begin{array}{ll}
\mathbb{R}: & 4^0 = 1 \\
\mathbb{N}_4\mathbb{C}: & 4^1 = 1+1+1+1 \\
\mathbb{N}_4\mathbb{Q}: & 4^2 = 1+2+3+4+3+2+1 \\
\mathbb{N}_4\mathbb{O}: & 4^3 = 1+3+6+10+12+12+10+6+3+1 \\
\mathbb{N}_4\mathbb{S}: & 4^4 = 1+4+10+20+31+40+44+40+31+20+10+4+1
\end{array}
\tag{50}
$$

In the last lines one can see the sedenions which do not produce division algebra. For both cases we have the unit element e_0 and the n basis elements:

$$\begin{aligned}
&\mathbb{R} \to \mathbb{TC} \to \mathbb{TQ} \to \mathbb{T}O \to \mathbb{T}S \\
&\mathbb{R} \to \mathbb{N}_4\mathbb{C} \to \mathbb{N}_4\mathbb{Q} \to \mathbb{N}_4\mathbb{O} \to \mathbb{N}_4\mathbb{S}
\end{aligned} \tag{51}$$

CONJUGATIONS CLASSESS AND ONE DIMENSIONAL REPRESENTATIONS

C_2	1	i	
$R^{(1)}$	1	1	z
$R^{(2)}$	1	-1	$\bar{z}$

.

The cyclic group C_3 has three conjugation classes, q_0, q and q^2, and, respectively, three one dimensional irreducible representations, $R^{(i)}$, $i = 1, 2, 3$. We write down the table of their characters, $\xi_l^{(i)}$:

$$\left(\begin{array}{c|ccc} - & 1 & q & q^2 \\ \hline \xi^{(1)} & 1 & 1 & 1 \\ \xi^{2)} & 1 & j_3 & j_3^2 \\ \xi^{(3)} & 1 & j_3^2 & j_3 \end{array}\right) \tag{68}$$

for C_3 ($j_3 =\equiv j = \exp\{2\pi \mathbf{i}/3\}$).

CONJUGATIONS CLASSESS AND ONE DIMENSIONAL REPRESENTATIONS

C_2	1	i	
$R^{(1)}$	1	1	z
$R^{(2)}$	1	-1	$\bar{z}$

.

The cyclic group C_3 has three conjugation classes, q_0, q and q^2, and, respectively, three one dimensional irreducible representations, $R^{(i)}$, $i = 1, 2, 3$. We write down the table of their characters, $\xi_l^{(i)}$:

$$\left(\begin{array}{c|ccc} - & 1 & q & q^2 \\ \hline \xi^{(1)} & 1 & 1 & 1 \\ \xi^{2)} & 1 & j_3 & j_3^2 \\ \xi^{(3)} & 1 & j_3^2 & j_3 \end{array}\right) \tag{68}$$

for C_3 ($j_3 =\equiv j = \exp\{2\pi \mathbf{i}/3\}$).

Ternary hyper-numbers

One can introduce the following basis forms:

$$
\begin{aligned}
K &= \frac{1}{3}(1+q+q^2), \\
E &= \frac{1}{3}(2-q-q^2), \\
I &= \frac{1}{\sqrt{3}}(q-q^2).
\end{aligned}
\tag{74}
$$

Reversely

$$
\begin{aligned}
q_0 &= K+E, \\
q &= K-\frac{1}{2}E+\frac{\sqrt{3}}{2}I, \\
q^2 &= K-\frac{1}{2}E-\frac{\sqrt{3}}{2}I.
\end{aligned}
\tag{75}
$$

Ternary hyper-numbers

The surface

$$x_0^3 + x_1^3 + x_2^3 - 3x_0x_1x_2 = \rho^3 \tag{82}$$

(see figure **??**) is a ternary analogue of the S^1 circle and it is related with the ternary Abelian group, $TU(1)$.

From the above figure one can see, that this surface approaches asymptotically the plane $x_0+x_1+x_2 = 0$ and the line $x_0 = x_1 = x_2$ orthogonal to it. In $\mathbb{T}_3\mathbb{C}$ they correspond to the ideals I_2 and I_1, respectively. The latter line will be called the "trisectrice".

One can compare this cubic surface to the quadratic surface -cylinder- what can be consider as equation $x_1^2 + x_2^2 = 1$ in the $\mathbb{C} \otimes \mathbb{R}$-space. This fact could be accepted as the first success of the ternary complex numbers of binary complex numbers what can be embedded only into even dimensional $\mathbb{R}^{2n}$ space.

$q_a^3 = q_0 \qquad a = 1,2,\ldots,8 \qquad j = \exp(\tfrac{2}{3}\pi i)$



$q_7\, q_1 = q_2$
$q_7\, q_2 = q_3$
$q_7\, q_3 = q_1$

$q_1\, q_7 = j\, q_2$
$q_2\, q_7 = j\, q_3$
$q_3\, q_7 = j\, q_1$

$q_1^2 = q_4$
$q_2^2 = q_5$
$q_3^2 = q_6$
$q_7^2 = q_8$

$q_1\, q_2\, q_3 = j^2\, q_0 \qquad q_2\, q_1\, q_3 = j\, q_0$

$q_4\, q_5\, q_6 = j^2\, q_0 \qquad q_5\, q_4\, q_6 = j\, q_0$

TOWARDS THE D+5,6- DIMENSIONAL EXTENSION OF LORENTZ GROUP

$$(x_0^{\,2} - x_1^{\,2} - x_2^{\,2} - x_3^{\,2})\, f(x_5, \ldots)$$

$$(x_0^{\,2} - x_1^{\,2} - x_2^{\,2} - x_3^{\,2})(x_5^{\,2} - x_6^{\,2})$$

R^n-COMPLEXIFICATION WITH FINITE GROUPS

Abelian Cyclic C _n- groups and Non-Abelian Groups

The Q_k operators with $k = 0, 1, 2, ..., 8$ satisfy to the following ternary S_3 commutation relations:

N	$\{klm\} \to \{n\}$	f^n_{klm}	N	$\{klm\} \to \{n\}$	f^n_{klm}	N	$\{klm\} \to \{n\}$	f^n_{klm}
1	$\{0,1,2\} \to \{6\}$	$\frac{1}{\sqrt{3}}$	29	$\{1,2,3\} \to \{0\}$	$\sqrt{3}$	57	$\{2,4,7\} \to \emptyset$	0
2	$\{0,1,3\} \to \{5\}$	$-\frac{1}{\sqrt{3}}$	30	$\{1,2,4\} \to \{2\}$	1	58	$\{2,4,8\} \to \emptyset$	0
3	$\{0,1,4\} \to \{0,7,8\}$	$\left\{0, 0, \sqrt{\frac{2}{3}}\right\}$	31	$\{1,2,5\} \to \{1\}$	1	59	$\{2,5,6\} \to \{6\}$	-1
4	$\{0,1,5\} \to \emptyset$	0	32	$\{1,2,6\} \to \emptyset$	0	60	$\{2,5,7\} \to \{0,7,8\}$	$\left\{\frac{3}{\sqrt{2}}, -1, -\frac{2}{\sqrt{3}}\right\}$
5	$\{0,1,6\} \to \emptyset$	0	33	$\{1,2,7\} \to \{6\}$	$-\sqrt{\frac{2}{3}}$	61	$\{2,5,8\} \to \{0,7,8\}$	$\left\{-\sqrt{\frac{3}{2}}, 0, 1\right\}$
6	$\{0,1,7\} \to \emptyset$	0	34	$\{1,2,8\} \to \{6\}$	$\sqrt{2}$	62	$\{2,6,7\} \to \emptyset$	0
7	$\{0,1,8\} \to \{1\}$	$-\sqrt{\frac{2}{3}}$	35	$\{1,3,4\} \to \{3\}$	-1	63	$\{2,6,8\} \to \emptyset$	0
8	$\{0,2,3\} \to \{4\}$	$\frac{1}{\sqrt{3}}$	36	$\{1,3,5\} \to \emptyset$	0	64	$\{2,7,8\} \to \{2\}$	$\frac{1}{\sqrt{3}}$
9	$\{0,2,4\} \to \emptyset$	0	37	$\{1,3,6\} \to \{1\}$	-1	65	$\{3,4,5\} \to \emptyset$	0
10	$\{0,2,5\} \to \{0,7,8\}$	$\left\{0, \frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{6}}\right\}$	38	$\{1,3,7\} \to \{5\}$	$\sqrt{\frac{2}{3}}$	66	$\{3,4,6\} \to \{4\}$	1
11	$\{0,2,6\} \to \emptyset$	0	39	$\{1,3,8\} \to \{5\}$	$\sqrt{2}$	67	$\{3,4,7\} \to \emptyset$	0
12	$\{0,2,7\} \to \{2\}$	$-\frac{1}{\sqrt{2}}$	40	$\{1,4,5\} \to \{5\}$	-1	68	$\{3,4,8\} \to \emptyset$	0
13	$\{0,2,8\} \to \{2\}$	$\frac{1}{\sqrt{6}}$	41	$\{1,4,6\} \to \{6\}$	1	69	$\{3,5,6\} \to \{5\}$	-1
14	$\{0,3,4\} \to \emptyset$	0	42	$\{1,4,7\} \to \{0,7,8\}$	$\left\{0, 0, \frac{1}{\sqrt{3}}\right\}$	70	$\{3,5,7\} \to \emptyset$	0
15	$\{0,3,5\} \to \emptyset$	0	43	$\{1,4,8\} \to \{0,7,8\}$	$\left\{\sqrt{6}, \sqrt{3}, 0\right\}$	71	$\{3,5,8\} \to \emptyset$	0
16	$\{0,3,6\} \to \{0,7,8\}$	$\left\{0, -\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{6}}\right\}$	44	$\{1,5,6\} \to \emptyset$	0	72	$\{3,6,7\} \to \{0,7,8\}$	$\left\{-\frac{3}{\sqrt{2}}, 1, -\frac{2}{\sqrt{3}}\right\}$
17	$\{0,3,7\} \to \{3\}$	$\frac{1}{\sqrt{2}}$	45	$\{1,5,7\} \to \emptyset$	0	73	$\{3,6,8\} \to \{0,7,8\}$	$\left\{-\sqrt{\frac{3}{2}}, 0, -1\right\}$
18	$\{0,3,8\} \to \{3\}$	$\frac{1}{\sqrt{6}}$	46	$\{1,5,8\} \to \emptyset$	0	74	$\{3,7,8\} \to \{3\}$	$\frac{1}{\sqrt{3}}$
19	$\{0,4,5\} \to \{3\}$	$-\frac{1}{\sqrt{3}}$	47	$\{1,6,7\} \to \emptyset$	0	75	$\{4,5,6\} \to \{0\}$	$-\sqrt{3}$
20	$\{0,4,6\} \to \{2\}$	$\frac{1}{\sqrt{3}}$	48	$\{1,6,8\} \to \emptyset$	0	76	$\{4,5,7\} \to \{3\}$	$\sqrt{\frac{2}{3}}$
21	$\{0,4,7\} \to \emptyset$	0	49	$\{1,7,8\} \to \{1\}$	$\frac{1}{\sqrt{3}}$	77	$\{4,5,8\} \to \{3\}$	$-\sqrt{2}$
22	$\{0,4,8\} \to \{4\}$	$\sqrt{\frac{2}{3}}$	50	$\{2,3,4\} \to \emptyset$	0	78	$\{4,6,7\} \to \{2\}$	$-\sqrt{\frac{2}{3}}$
23	$\{0,5,6\} \to \{1\}$	$-\frac{1}{\sqrt{3}}$	51	$\{2,3,5\} \to \{3\}$	1	79	$\{4,6,8\} \to \{2\}$	$-\sqrt{2}$
24	$\{0,5,7\} \to \{5\}$	$\frac{1}{\sqrt{2}}$	52	$\{2,3,6\} \to \{2\}$	1	80	$\{4,7,8\} \to \{4\}$	$-\frac{1}{\sqrt{3}}$
25	$\{0,5,8\} \to \{5\}$	$-\frac{1}{\sqrt{6}}$	53	$\{2,3,7\} \to \{4\}$	$2\sqrt{\frac{2}{3}}$	81	$\{5,6,7\} \to \{1\}$	$-2\sqrt{\frac{2}{3}}$
26	$\{0,6,7\} \to \{6\}$	$-\frac{1}{\sqrt{2}}$	54	$\{2,3,8\} \to \emptyset$	0	82	$\{5,6,8\} \to \emptyset$	0
27	$\{0,6,8\} \to \{6\}$	$-\frac{1}{\sqrt{6}}$	55	$\{2,4,5\} \to \{4\}$	-1	83	$\{5,7,8\} \to \{5\}$	$-\frac{1}{\sqrt{3}}$
28	$\{0,7,8\} \to \emptyset$	0	56	$\{2,4,6\} \to \emptyset$	0	84	$\{6,7,8\} \to \{6\}$	$-\frac{1}{\sqrt{3}}$

We have got 84 commutations relations. One commutation relation, $\{Q_0, Q_7, Q_8\}_{S_3} = 0$, provide the Cartan subalgebra. Let separate the rest 83 commutation relations on the 5 groups$(18+18+27+18+2)$. The first group contains itself the following 18 commutation

ТЕМЫ

ДЛЯ НАУЧНО-ПРАКТИЧЕСКИХ РАБОТ СТУДЕНТОВ И АСПИРАНТОВ ,

УЧИТЫВАЮЩИХ НАУЧНЫЕ НАПРАВЛЕНИЯ ИССЛЕДОВАНИЙ. ПРОВОДИМЫХ В НИУ-6

1.ПРАМАТЕРИЯ И ОБРАЗОВАНИЕ ВСЕЛЕННОЙ.

НЕСТАБИЛЬНОСТЬ ПРОТОНА И ПОСТ – ЯДЕРНАЯ ЭНЕРГЕТИКА .

ЦВЕТОВАЯ КВАРКОВАЯ МОДЕЛЬ НУКЛОНОВ – ДОСТИЖЕНИЯ И ПРОБЛЕМЫ ЗАПИРАНИЯ ЦВЕТА

ПРОБЛЕМЫ ЯДЕРНОЙ ФИЗИКИ - КВАНТОВАНИЕ НУКЛОНОВ – ПРОТОН + НЕЙТРОН .

ПРОБЛЕМЫ ТРОЙСТВЕННОСТИ МАТЕРИИ. НОВЫЕ ВИДЫ НЕСТАБИЛЬНОЙ МАССИВНОЙ МАТЕРИИ.

СТРУКТУРА ЭЛЕКТРОНА.

ПРАМАТЕРИЯ И ЕДИНАЯ МОДЕЛЬ НУКЛОНОВ И ЭЛЕКТРОНА (P,N,E).

РОЛЬ ПРАМАТЕРИИ В ПРОЦЕССАХ ОБРАЗОВАНИЯ ЭЛЕКТРОМАГНИТНО -ЗАРЯЖЕННОЙ

ЯДЕРНОЙ И ЛЕПТОННОЙ МАТЕРИИ.

ПРОБЛЕМА АНТИ - ВЕЩЕСТВА.

ПРОБЛЕМЫ ПОСТРОЕНИЯ ГЕОМЕТРИИ ВСЕЛЕННОЙ: ДОПОЛНИТЕЛЬНЫЕ ПРОСТРАНСТВЕННЫЕ

ИЗМЕРЕНИЯ И ВТОРОЕ ВРЕМЯ

РАСШИРЕНИЕ ТЕОРИИ ОТНОСИТЕЛЬНОСТИ И НОВЫЕ ВИДЫ ЭНЕРГИИ.

ТЕОРИЯ И ПРАКТИЧЕСКИЕ ПРИМЕНЕНИЯ.

- of light in Minkowsky ($D = 3 + 1$) space-time. Absence of singularities in such a spacetime
- allows you to enter the gauge invariance in a region, which can connect two kinds of
- matter: the matter substance and radiation. The substance described by the fundamental
- fermion fields with spin 1/2[4], G.Weyl,[11], and radiation - gauge fields with spin 1. The
- question of maintaining gauge invariance may depend on the existence of singularities in
- this space-time, which can be a source of symmetry breaking. This option is actually
- a violation of gauge symmetry associated with the existence of space-time singularities
- at small or large distances. Note that the existence of singularities at small distances
- can lead to a change of the Riemann metric and, therefore, to a dynamical violation of
- space-time Lorentz symmetry (see for example, [7]).

- Thus the formalism of quantum field theory includes the geometric foundation
- of space-time picture of the ”visible ”world and the operator-functional methods of describing
- a matter moving and interacting in this environment. But now some phenomena
- in physics of elementary particles pose the question the need to expand our notions of
- space and time?! In this case the first question arises of dimension and signature of a
- new hypothetical world. In our opinion, now modern science close to understanding to
- the origin of the visible part of universe defined by a D=(3+1)-dimensional space-time
- continuum, obeying to the laws of absolutism speed of light, and the observable fermion
- matter of which has the ”unified” electromagnetic nature. In articles [2] , [19] it was
- suggested that only the Dirac fermion matter can satisfy to the laws of absolutism speed

LIE algebras SO(3,1) and SL(2C)

- The Lie algebra of Lorentz group $SO(3, 1)$ is isomorphic to the algebra of its double covering
- ***Spin*(3, 1) = *SL*(2*C*)-**group
- the irreducible representations of what can
- be defined by two integer or semi-integer numbers (μ, v) of the finite-dimensional representations of the $SU(2) \times SU(2)$ group.
- The minimal representations of this group are
- Scalar (0, 0) representation
- Weyl spinors , (1/2, 0)L- and (0, 1/2)R-representations
- what are related by P -parity operation (and complex conjugation):
- $x0 \rightarrow x0, \ \vec{x} \rightarrow -\vec{x}$, $(1/2, 0) \rightarrow (0, 1/2)$.

НАЧАЛО SU(2)_SWxU(1)_Y

- To describe the *I*_- charged weak currents
- and combine them to the EM- currents in Weinberg-Salam model $SU(2)WI \times U(1)Y$ it
- was used the ideas of the Heisenberg $SU(2)I$ - isotopic group and the following relation
- $$Q(EM) = I_3 + Y/2.$$

- As one of the main result of a such model it was predicted the neutral
- weak interactions what was experimentally confirmed in GARGAMELLE CERN neutrino
- experiments in 1974 year. In this model the P- violation (C-violation) was constructed by
- hands taking the left- and right handed fermions in different $SU(2)WI$ - representations.
- For breaking the gauge symmetry in Weinberg- Salam model it was used the mechanism
- in the internal sector of the model what predicted the existence of a new fundamental
- scalar particle- Higgs boson.

- . Mechanism of the appearance of
- the masses of gauge bosons and fermions is enough formal and it is not clear its link to
- structural changes of the space-time. At least , in spite of preliminary of strong indications
- and a lot of discussions CERN plans to continue these experiments for the future cycle
- of LHC-collider work with planing to get much more the energy of the proton beams.
- Fermilab also resumed the work on the improvement of the Tevatron to finally clarify the

- nature of signals detected at CERN collider at energy of 125 Gev. One can propose that
- the role of the weak sector of the Standard Model is the way to understand the origin
- of the visible universe. More suppressed processes going to the CP- violation may be
- associated with an unknown dynamics at the more smaller distances $\sim 10^{-16-17}$cm. In
- addressing this issue again we faced with the dilemma of the mechanism of these phenomena:
- defects of the space-time geometry or/and a new dynamics related to the new
- interactions ? Obviously the issue is closely related to another important problem - the
- existence of three quark-lepton families. All experimental information on three family
- mixing and CP-violation can be encoded into the Cabibbo-Kobajashi-Maskawa (CKM)-
- matrix parameters which also requires explanation!

- The 3- family mixing explanation is
- completely going into the mass origin problem. In the second case one should again to
- study the problem related to the local gauge symmetry breaking - ”Higgsology” or unknown
- a space-time singularity structure. In the depths of this phenomenology is waiting
- for us very rich physics what can shed light on the production the visible part of Universe!

Towards a new spinor-fermion structure

- we do not define the fermion matter that fills the space-time continuum should have a universal property, *i.e.* Dirac half-one fermions[2].
- [GV],[AV].
- It means that we can imagine the existence of exotic fermion matter , for example,
- having another spin $1/n$, $n \geq 3$ and without an electromagnetic (color) charge
- nature. In this picture our visible Dirac Universe forming a topological cycle could be
- embedded into Meta - Universe having much more reach the space-time topology

- . As a
- messenger between the cycles we suggested neutrinos with Dirac mass equal to zero.
- To construct the spin 1*/n* fermion theories first one should find out the examples of
- the geometrical spaces having such a spinor structure [19],[20],[22]. Taking into account
- a possibility to imply the spaces with arbitrary spin structure in formulation of the basic
- principles of the string theory one could significantly expand the assumption touching
- the D- dimensional pseudo-Lorentz space, in which the string is moving . We think that
- in this case the string and superstrings theories could considerably extend the set of
- predictions for modern physics of elementary particles. In according to such geometrical
- objects one can search for new symmetries, what we already started to study in the class
- of n-ary symmetries with corresponding n-ary algebras what already have been discussed
- in literature, for example, [6],[22], and reference there.

РАСШИРЕНИЕ СИММЕТРИИ ЛОРЕНЦА.

-
- $[S_i, S_j] = i\varepsilon_{ijk} S_{k,}\ \sigma_i^2 = \hat{1}_2 = \sigma_0$
- $S_j = \frac{1}{2}\sigma_j, j = 1,2,3$
- $\sigma_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma_1 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$
- $[Q_a, Q_b] = 0, Q_a^3 = \hat{1}_3 = Q_0$
-
- $Q_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, Q_1 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, Q_2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$

GROUP ALGEBRAS -GEOMETRY

- $\Lambda_{ia} = S_i \otimes Q_a = \left(\frac{1}{2}\sigma_i\right) \otimes Q_a,$ i=0,1,2,3;a=0,1,2
- $\hat{\Lambda} = \left\{\sum_{k=0}^{k=3} x_k S_k\right\} \otimes \left\{\sum_{c=0}^{2} y_c Q_c\right\}$
-
- $\hat{\Lambda} =$

$$\frac{1}{2}\begin{pmatrix} (x_0+x_3)y_0 & (x_0+x_3)y_1 & (x_0+x_3)y_2 & (x_1-ix_2)y_0 & (x_1-ix_2)y_1 & (x_1-ix_2)y_2 \\ (x_0+x_3)y_2 & (x_0+x_3)y_0 & (x_0+x_3)y_1 & (x_1-ix_2)y_2 & (x_1-ix_2)y_0 & (x_1-ix_2)y_1 \\ (x_0+x_3)y_1 & (x_0+x_3)y_2 & (x_0+x_3)y_0 & (x_1-ix_2)y_1 & (x_1-ix_2)y_2 & (x_1-ix_2)y_0 \\ (x_1+ix_2)y_0 & (x_1+ix_2)y_1 & (x_1+ix_2)y_2 & (x_0-x_3)y_0 & (x_0-x_3)y_1 & (x_0-x_3)y_2 \\ (x_1+ix_2)y_2 & (x_1+ix_2)y_0 & (x_1+ix_2)y_1 & (x_0-x_3)y_2 & (x_0-x_3)y_0 & (x_0-x_3)y_1 \\ (x_1+ix_2)y_1 & (x_1+ix_2)y_2 & (x_1+ix_2)y_0 & (x_0-x_3)y_1 & (x_0-x_3)y_2 & (x_0-x_3)y_0 \end{pmatrix}$$

GROUP ALGEBRAS -GEOMETRY

- $\Lambda_{ai}^{T} = Q_a \otimes S_i = Q_a \otimes \left(\frac{1}{2}\sigma_i\right),$
  i=0,1,2,3;a=0,1,2
- $Q_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, Q_1 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, Q_2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$
- $\hat{\Lambda}^{T} = \{\sum_{c=0}^{2} y_c Q_c\} \otimes \{\sum_{k=0}^{k=3} x_k S_k\}$
- $= \{\sum_{c=0}^{2} y_c Q_c\} \otimes \left\{\frac{1}{2}\sum_{i=0}^{3} x_i \sigma_i\right\}$

GROUP ALGEBRAS -GEOMETRY

- $\Lambda_{ai}^{T} = Q_a \otimes S_i = Q_a \otimes \left(\frac{1}{2}\sigma_i\right)$,
  i=0,1,2,3;a=0,1,2
- $Q_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, Q_1 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, Q_2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$
- $\hat{\Lambda}^{T} = \{\sum_{c=0}^{2} y_c Q_c\} \otimes \{\sum_{k=0}^{k=3} x_k S_k\} =$
  $\{\sum_{c=0}^{2} y_c Q_c\} \otimes \left\{\frac{1}{2}\sum_{i=0}^{3} x_i \sigma_i\right\}$

GROUP ALGEBRAS -GEOMETRY

-
- $\hat{\Lambda}^T(SU(2Q) = \frac{1}{2}\begin{pmatrix} y_0 & y_1 & y_2 \\ y_2 & y_0 & y_1 \\ y_1 & y_2 & y_0 \end{pmatrix} \otimes \begin{pmatrix} x3 & x_1 - ix_2 \\ x_1 + ix_2 & -x_3 \end{pmatrix} =$
-
- $= \frac{1}{2}\begin{pmatrix} x_3y_0 & (x_1-ix_2)y_0 & x_3y_1 & (x_1-ix_2)y_1 & x_3y_2 & (x_1-ix_2)y_2 \\ (x_1+ix_2)y_0 & -x_3y_0 & (x_1+ix_2)y_1 & -x_3y_1 & (x_1+ix_2)y_2 & -x_3y_2 \\ x_3y_2 & (x_1-ix_2)y_2 & x_3y_0 & (x_1-ix_2)y_0 & x_3y_1 & (x_1-ix_2)y_1 \\ (x_1+ix_2)y_2 & -x_3y_2 & (x_1+ix_2)y_0 & -x_3y_0 & (x_1+ix_2)y_1 & -x_3y_1 \\ x_3y_1 & (x_1-ix_2)y_1 & x_3y_2 & (x_1-ix_2)y_2 & x_3y_0 & (x_1-ix_2)y_0 \\ (x_1+ix_2)y_1 & -x_3y_1 & (x_1+ix_2)y_2 & -x_3y_2 & (x_1+ix_2)y_0 & -x_3y_0 \end{pmatrix}$

SU(2Q)-ALGEBRA

- $\hat{\Lambda} = \left\{\sum_{k=0}^{k=3} x_k S_k\right\} \otimes \left\{\sum_{c=0}^{2} y_c Q_c\right\}$
-
- $\left[\Lambda_{ia}, \Lambda_{jb}\right] = i\varepsilon_{ijk}\eta_c \Lambda_{kc},$
- $\Lambda_{kc} = S_k \otimes Q_c, Q_c = Q_a Q_b = \eta_{abc} Q_c$
- $\left[S_i \otimes Q_a, S_j \otimes Q_b\right] = S_i \cdot S_j \otimes Q_a \cdot Q_b - S_j \cdot S_i \otimes Q_b \cdot Q_a = \left(S_i \cdot S_j - S_j \cdot S_i\right) \otimes Q_a Q_b =$
- $= i\varepsilon_{ijk} S_k \otimes \eta_{abc} Q_c = i\varepsilon_{ijk} S_k \otimes Q_a Q_b = i\varepsilon_{ijk} S_k \otimes \eta_{abc} Q_c$

SU(2Q)-ALGEBRA

- $\hat{\Lambda}(SU(2Q)) =$

$$\frac{1}{2}\begin{pmatrix} x_3y_0 & x_3y_1 & x_3y_2 & (x_1-ix_2)y_0 & (x_1-ix_2)y_1 & (x_1-ix_2)y_2 \\ x_3y_2 & x_3y_0 & x_3y_1 & (x_1-ix_2)y_2 & (x_1-ix_2)y_0 & (x_1-ix_2)y_1 \\ x_3y_1 & x_3y_2 & x_3y_0 & (x_1-ix_2)y_1 & (x_1-ix_2)y_2 & (x_1-ix_2)y_0 \\ (x_1+ix_2)y_0 & (x_1+ix_2)y_1 & (x_1+ix_2)y_2 & -x_3y_0 & -x_3y_1 & -x_3y_2 \\ (x_1+ix_2)y_2 & (x_1+ix_2)y_0 & (x_1+ix_2)y_1 & -x_3y_2 & -x_3y_0 & -x_3y_1 \\ (x_1+ix_2)y_1 & (x_1+ix_2)y_2 & (x_1+ix_2)y_0 & -x_3y_1 & -x_3y_2 & -x_3y_0 \end{pmatrix}$$

-

SU(3^c)XU(1)XSU(2)XU(1)xSU(3H)XU(1H)
4-dimsuperstrings
With World-Sheet Fermions(1992Padova)

Table 2: **The list of quantum numbers of the states. Model 1.**

N^o	$b_1, b_2, b_3, b_4, b_5, b_6$	SO_{hid}	$U(5)^I$	$U(3)^I$	$U(5)^{II}$	$U(3)^{II}$	Y_5^I	Y_3^I	Y_5^{II}	Y_3^{II}
1	RNS		5	3	1	1	–1	–1	0	0
			1	1	5	$\bar{3}$	0	0	–1	–1
	0 2 0 1 2(6) 0		5	1	5	1	–1	0	–1	0
$\hat{\Phi}$			1	3	1	3	0	1	0	1
			5	1	1	3	–1	0	0	1
			1	3	5	1	0	1	–1	0
2	0 1 0 0 0 0		1	3	1	1	5/2	–1/2	0	0
			$\bar{5}$	3	1	1	–3/2	–1/2	0	0
			10	1	1	1	1/2	3/2	0	0
$\hat{\Psi}$	0 3 0 0 0 0		1	1	1	1	5/2	3/2	0	0
			$\bar{5}$	1	1	1	–3/2	3/2	0	0
			10	3	1	1	1/2	–1/2	0	0
3	0 0 1 1 3 0	$-_1\ \pm_2$	1	1	1	3	0	–3/2	0	–1/2
	0 0 1 1 7 0	$-_1\ \pm_2$	1	$\bar{3}$	1	1	0	1/2	0	3/2
$\hat{\Psi}^H$	0 2 1 1 3 0	$+_1\ \pm_2$	1	$\bar{3}$	1	3	0	1/2	0	–1/2
	0 2 1 1 7 0	$+_1\ \pm_2$	1	1	1	1	0	–3/2	0	3/2
4	1 1 1 0 1 1	$\mp_1\ \pm_3$	1	1	1	$\bar{3}$	0	–3/2	0	1/2
	1 1 1 0 5 1	$\mp_1\ \pm_3$	1	$\bar{3}$	1	1	0	1/2	0	–3/2
$\hat{\Phi}^H$	1 3 1 0 1 1	$\pm_1\ \pm_3$	1	$\bar{3}$	1	$\bar{3}$	0	1/2	0	1/2
	1 3 1 0 5 1	$\pm_1\ \pm_3$	1	1	1	1	0	–3/2	0	–3/2
5	0 1(3) 1 0 2(6) 1	$-_1\ \pm_3$	1	3($\bar{3}$)	1	1	$\pm 5/4$	$\pm 1/4$	$\pm 5/4$	$\mp 3/4$
		$+_1\ \pm_3$	5($\bar{5}$)	1	1	1	$\pm 1/4$	$\mp 3/4$	$\pm 5/4$	$\mp 3/4$
$\hat{\phi}$	0 1(3) 1 0 4 1	$-_1\ \pm_3$	1	1	1	3($\bar{3}$)	$\pm 5/4$	$\mp 3/4$	$\pm 5/4$	$\pm 1/4$
		$+_1\ \pm_3$	1	1	5($\bar{5}$)	1	$\pm 5/4$	$\mp 3/4$	$\pm 1/4$	$\mp 3/4$
6	1 2 0 0 3(5) 1	$\pm_1\ -_4$	1	1	1	1	$\pm 5/4$	$\pm 3/4$	$\mp 5/4$	$\mp 3/4$
	1 1(3) 0 1 5(3) 1	$+_1\ \mp_4$	1	1	1	1	$\pm 5/4$	$\pm 3/4$	$\pm 5/4$	$\pm 3/4$
$\hat{\sigma}$	0 0 1 0 2(6) 0	$\mp_3\ +_4$	1	1	1	1	$\pm 5/4$	$\mp 3/4$	$\pm 5/4$	$\mp 3/4$

1999-2000-Padova-CERN

Can Neutrinos Probe Extra Dimensions?

V. Ammosov[b] and G. Volkov[a, b]

[a] *Universita di Padova, Dipartimento di Fisica Galileo Galilei, INFN, Padova , Italy.*

[b] *Institute for High Energy Physics, Protvino, Russia.*

СВОЙСТВА НЕЙТРИНО О МНОГОМЕРНОМ ОБОБЩЕНИИ ТЕОРИИ ОТНОСИТЕЛЬНОСТИ

ПРЕДСТАВЛЕНИЯ ГРУППЫ ЛОРЕНЦ- ПУАНКАРЕ

- *1. Спин s=½*
- ***2. Майорано-Вейлевская природа** - 2-х компклексные степени свободы*

3. Масса m =O(eV)

ЭЛЕКТРОМАГНИТНАЯ СТЕРИЛЬНОСТЬ

- *4. Заряд Q (EM)= O*
- *5. Магнитный момент Mag=O(0)*

(V – A)- СЛАБАЯ СВЯЗЬ С ЗЛЕКТРОМАГНИТНЫМ МИРОМ

- ***6. Взаимодействие слабое***

Letter of Intent

Measurement of Neutrino Velocities at the CERN WANF using Bare Target Neutrino Beams

F. Dydak, J. Ellis, V. Falaleev, J. Gareyte
CERN, CH-1211, Geneva 23, Switzerland

V. Ammosov, G. Gapienko, G. Volkov
Institute for High Energy Physics
RU-142284 Protvino, Moscow region, Russia

Abstract

String/D-branes theories might establish the connection between the geometrical properties of the gauge forces, the associated vacuum and the role of the "sterile" particles. It was pointed out that the speed of a "sterile" particle can be differ from standard expectations of the Special Theory of Relativity.

- **References**

- [1] A.N.Amaglobeli, A.R.Kereselidze, A.G.Liparteliani,G.G.Volkov, *The supersymmetric*
- *vector-like horizontal model with intermediate symmetry breaking scale Phys. Lett.* **B237**
- (1990) 417;
- 263
- Alexander Maslikov and Guennady Volkov
- [2] V. Ammosov, G. Volkov, *Can Neutrinos Probe Extra Dimensions?* hep-ph/0008032. Padua
- preprint DFPD-00/TH/39, arXiv:hep-ph/0008032v1 (2000).
- [3] D.S. Baranov and G.G.Volkov *Neutrino and Extra World.* arXiv:1211.4708
- *”Neutrino On The Possible New Time Structure”* arXiv:1302.1482

- [6] A.Dubrovskiy and G.Volkov, *Ternary numbers and algebras. Reflexive numbers and Berger*
- *graphs* Adv.Appl.CliffordAlgebras17:159-181,2007
- archiv:hep-th/0608073, (2006).
- [7] J. R. Ellis, N. E. Mavromatos, D. V. Nanopoulos and G. Volkov, *Gravitational-Recoil Effects*
- *on Fermion Propagation in Space-Time Foam* Gen. Rel. Grav. 32 (2000)1777 [arXiv:grqc/
- 9911055].
- [8] R.W. Hamilton, *On Quaternions*, Proceedings of the Royal Irish Academy, Nov.11 (1844),
- **v.3** (1847), 1-16.
- [9] L.N. Lipatov, A. Sabio-Vera, V.N. Velizhanin and G.G. Volkov, *From New Geometry Towards*
- *a New Symmetry. Reflexive Numbers and Berger Graphs from Calabi-Yau Spaces*,
- Dark Matter in Astro- and Particle Physics , **DARK-2004** Springer, (2004) 622-655, ISBN-
- 13 978-3-540-26372-2 Springer Berlin Heidelberg New York; Int.J.Mod.Phys. A21 (2006)
- 2953-3006
- [10] L. N. Lipatov, M. Rausch de Traunbenberg, G. Volkov, *On the ternary complex analysis*
- *and its applications* J. Math. Phys. **49** 013502 (2008)

- [14] V. Monich, B. Struminsky, G. Volkov, *Oscillation and CP violation in the "horizontal"*
- *superweak gauge scheme* Phys. Lett.**B104** p. 382-384(1981)
- [15] H. Poincarŕe *Sur les propriŕetŕes du potential et sur les fonctions Abeliennes* Acta Matematica
- **22**(1) 89-178.
- [16] V. Samoylenko and G.Volkov *The GUT of the light: On the Abelian Complexifications of*
- *the Euclidean Rn spaces* arXiv:0912.2037 (2009)
- [17] G.G. Volkov, A.G. Liparteliani, V.A. Monich, Y.P. Nikitin, *Mixing of leptons with different*
- *quantum numbers in gauge schemes of the weak and electromagnetic interactions* Sov. J.
- Nucl. Phys. (Engl. Transl.); **27**(1978)735
- Yad.Fiz. **27** (1978) 1395-1402, preprint IFVE-77-89.
- [18] G. Volkov, *Hunting for the New Symmetries in Calabi-Yau Jungles*, Int.J. Mod. Phys **A19**
- (2004) 4835-4860, hep-th/0402042.
- [19] G.G. Volkov, Annales Fond. Broiglie **31** (2006) 227
- *Geometry of Majorana neutrino and new symmetries*
- *short version in* hep-ph/0607334.
- [20] G. Volkov, *The possible signals from D=6* arxiv:1112.3583(2011)
- [21] G. Volkov *On the complexifications of the Euclidean Rn spaces and the n-dimensional*
- *generalization of Pythagore theorem* arXiv:1006.5630(2010)
- [22] G. Volkov *Ternary "Quaternions" and Ternary TU*(3) *algebra* arXiv:1006.5627 (2010)
- 264